

doi <https://doi.org/10.61186/nl.3.2.1>

# Level of CSF GAP-43 and white matter microstructural changes in Alzheimer's disease

Marjan Assefi<sup>1\*</sup>, Alireza Sharafshah<sup>2</sup>, Atefeh Ashtari<sup>1</sup>, Sayeh Afshar<sup>1</sup>, Keysan Pour Moghtader<sup>2</sup>, Yasir Waheed<sup>3</sup>, and the Alzheimer's Disease Neuroimaging Initiative\*\*

- 1- Marie Curie Science Research Center, Greensboro, USA
- 2- Department of Biology, Marie Curie Science Research Center, Greensboro, USA
- 3- MEU Research Unit, Middle East University, Amman, 11831, Jordan

Received 4 January 2024; Revised 22 March 2024; Accepted 29 March 2024; Published 29 April 2024

## Abstract

**Objectives:** Several studies have reported altered cerebrospinal fluid (CSF) concentrations of presynaptic proteins, such as growth-associated protein 43 (GAP-43) in Alzheimer's disease (AD) patients. Given the potential predictive role of CSF GAP-43 for AD, the current study aimed to investigate the relationship between CSF GAP-43 levels and DTI-detected microstructural changes in the white matter (WM).

**Methods:** Data from three groups of participants including 39 control normals (CN), 138 MCI, and 39 AD subjects were obtained from the Alzheimer's disease Neuroimaging Initiative (ADNI). Linear regression was used to the association of CSF-GAP43 and DTI values (including MD, RD, AxD, and FA) in the brain.

**Results:** We found a significant association between CSF-GAP43 and FA ( $p$ -value = 0.011). Also, a negative association was found between CSF-GAP43 concentration and AD, MD, and RD values in MCI ( $p$ -value = 0.013,  $p$ -value = 0.004,  $p$ -value = 0.017). The regression models also revealed a positive association between CSF-GAP43 and FA value in AD subjects ( $p$ -value = 0.028). Furthermore, increased CSF-GAP43 level was associated with lower AD, MD, and RD values in brain WM of AD patients ( $p$ -value = 0.022,  $p$ -value = 0.033,  $p$ -value = 0.041).

**Conclusion:** Our study provides a better understanding of the link between CSF GAP-43 and WM changes in patients with MCI and AD. Our findings support the application of CSF GAP-43 as an effective biomarker for monitoring individuals at high risk of AD in the early stages.

**Keywords:** Alzheimer's Disease, white matter, GAP-43, synaptic, DTI

\*Correspondence to Marjan Assefi, Marie Curie Science Research Center, Greensboro, USA

Email: [massefi@aggies.ncat.edu](mailto:massefi@aggies.ncat.edu)

\*\*Data used in the preparation of this article were obtained from the Alzheimer's disease Neuroimaging Initiative (ADNI) database ([adni.loni.usc.edu](http://adni.loni.usc.edu)). As such, the investigators within the ADNI contributed to the design and implementation of ADNI and/or provided data but did not participate in the analysis or writing of this report. A complete listing of ADNI investigators can be found at: [http://adni.loni.usc.edu/wp-content/uploads/how\\_to\\_apply/ADNI\\_Acknowledgement\\_List.pdf](http://adni.loni.usc.edu/wp-content/uploads/how_to_apply/ADNI_Acknowledgement_List.pdf)



**Cite this article as:** Assefi, M., Sharafshah, A., Ashtari, A., Afshar, S., Pour Moghtader, K., Waheed, Y. Level of CSF GAP-43 and white matter microstructural changes in Alzheimer's disease. *Neurology Letters*, 2024; 3(Supplementary 1 (Diagnostic and Therapeutic advances in Neurodegenerative diseases)): 1-6. doi: 10.61186/nl.3.2.1.

## Introduction

Alzheimer's disease (AD) is one of the most common neurodegenerative disorders and the leading cause of dementia worldwide. The neuropathological hallmarks of AD include the extracellular aggregation of amyloid- $\beta$  ( $A\beta$ ) plaques, intracellular deposition of neurofibrillary tangles containing phosphorylated tau protein, and synaptic loss (1). Synapses are crucial for cognitive function, and synaptic impairment is a

pathological feature of AD. Quantitative morphometric studies have demonstrated a 25-35% decline in synaptic density in the temporal and frontal cortical biopsies of patients within 2-4 years of AD onset (2). In AD patients, synaptic loss primarily occurs in the neocortex and hippocampus (3). Synaptic damage and cell loss lead to brain atrophy, and synaptic dysfunction is associated with cognitive impairment in AD (4). Evaluating synaptic dysfunction in vivo could guide AD clinical research and provide biomarkers for outcome

**Table 1.** Participants characteristic

Demographic and health characteristics	CN (39)	MCI(138)	AD(39)	<i>P-value</i>
Age, years	73.1(±6.3)	72.8(±6.8)	74.4 (±8.1)	0.655
Education, years	16.4(±2.6)	16.0(±2.6)	15.2(±2.9)	0.084
MMSE score	28.8(±1.4)	27.9(±1.9)	23.4(±1.8)	<0.001
CSF GAP-43 level, pg/mL	4586.9 (±1240.4)	5322.7(±1539.6)	6083.3(±1136.3)	<0.001

Values are shown as mean(±SD), Mini-Mental State Examination(MMSE), Control normal (CN), Mild cognitive impairment (MCI), and Alzheimer's disease (AD); results of ANOVA analysis between groups noted as p-value

assessment in AD clinical trials. In recent years, progress has been made in the evaluation of synaptic biomarkers in the cerebrospinal fluid (CSF). Depending on the localization of the synaptic protein, synaptic biomarkers can be divided into pre- and postsynaptic biomarkers. Several studies have reported altered CSF concentrations of presynaptic proteins, such as growth-associated protein 43 (GAP-43), synaptosomal-associated protein 25 (SNAP-25), and synaptotagmin-1, as well as the postsynaptic protein neurogranin, in AD patients (5-8). Among these synaptic proteins, GAP-43 plays an important role in the learning and memory storage process (9, 10).

Diffusion Tensor Imaging (DTI) enables the in vivo, non-invasive assessment of neurodegeneration, white matter (WM) disruption, and synaptic damage in patients with Alzheimer's disease (AD) (10). Emerging research suggests that WM changes may serve as a marker for pathological significance, potentially offering a promising target for the early diagnosis of dementia (11). The evidence regarding the association between CSF GAP-43 and WM microstructure in the AD continuum is currently insufficient. Given the potential predictive role of CSF GAP-43 for AD, the current study aimed to investigate the relationship between CSF GAP-43 levels and DTI-detected microstructural changes in the WM.

## Materials and Methods

### Data Acquisition

The data for this investigation was obtained from the Alzheimer's Disease Neuroimaging Initiative (ADNI) database ([adni.loni.usc.edu](http://adni.loni.usc.edu)). ADNI was established in 2003 as a public-private partnership directed by Principal Investigator Michael W. Weiner, MD. The primary objective of ADNI is to assess the progression of mild cognitive impairment (MCI) and early Alzheimer's disease (AD) by combining serial PET, MRI, biological markers, and clinical and neuropsychological measures. The most up-to-date information is available at [www.adni-info.org](http://www.adni-info.org).

### Participants

The study included baseline data from participants in the ADNI-2 and ADNI-GO cohorts who had available CSF GAP-43 and DTI statistical results. The sample comprised 39 control normals (CN), 138 MCI, and 39 AD subjects. All MCI subjects were diagnosed based on the following criteria: Mini-Mental


State Examination (MMSE) scores between 24 and 30, a memory complaint, objective memory loss measured by education-adjusted scores on the Wechsler Memory Scale Logical Memory II, a Clinical Dementia Rating (CDR) of 0.5, absence of significant impairment in other cognitive domains, essentially preserved activities of daily living, and absence of dementia.

### CSF GAP-43 Measurement

The cerebrospinal fluid (CSF) levels of growth-associated protein 43 (GAP-43) were measured using an in-house enzyme-linked immunosorbent assay (ELISA) method developed at the Clinical Neurochemistry Lab, University of Gothenburg, Sweden. The ELISA was developed by using a combination of the monoclonal GAP-43 antibody NM4 as the coating antibody and the polyclonal GAP-43 antibody ABB-135 as the detector antibody, which recognizes the C-terminal of GAP-43. The ELISA assay range for CSF-GAP43 was from 312 to 20,000 pg/ml. Quality control CSF samples were provided by the Clinical Neurochemistry Laboratory, Sahlgrenska University Hospital, Mölndal, Sweden. The details of the ELISA assay for the measurement of CSF GAP-43 have been described previously.

### DTI Processing and Image Analysis

The results of the DTI regions of interest (ROI) analysis were downloaded from the ADNI cohort. DTI scans underwent normalization using the Montreal Neurological Institute and Hospital (MNI) `nu_correct` tool ([www.bic.mni.mcgill.ca/software/](http://www.bic.mni.mcgill.ca/software/)). Non-brain tissues were removed using the Brain Extraction Tool (BET) from FSL (12). The T1-weighted image was aligned to a version of the Colins27 brain template (13) using FSL's `flirt` (14). A single diffusion tensor was modeled at each voxel in the brain [19]. Scalar anisotropy and diffusivity maps were obtained from the resulting diffusion tensor eigenvalues ( $\lambda_1$ ,  $\lambda_2$ ,  $\lambda_3$ ). Fractional anisotropy (FA), mean diffusivity (MD), radial diffusivity (RD), and axial diffusivity (AxD) were calculated. Lower FA and higher RD, AxD, and MD are related to demyelination and degeneration in white matter. The FA image from the Johns Hopkins University (JHU) DTI atlas was registered to each subject using a shared information-based elastic registration algorithm. To prevent label intermixing, nearest-neighbor interpolation was used to apply the deformation to the stereotaxic JHU "Eve" white matter atlas labels. This placed the



**Figure 1.** Association between CSF-GAP43 and DTI measures in CN subjects.

atlas ROIs in the same coordinate space as the DTI maps. The average FA and MD were then calculated within the boundaries of each ROI mask for each subject. Tensor-based spatial statistics and tract-based spatial statistics (TBSS) were also performed to extract the mean FA in the ROIs along with the skeleton.

### Statistical Analysis

The SPSS software (Statistical Package for the Social Sciences, version 16, USA) was used for data analysis. First, simple linear regression models were conducted to assess the association of CSF-GAP43 with clinical and demographic. Next, the association of CSF-GAP43 and DTI values (including MD, RD, AxD, and FA) in the brain was measured using simple linear regression models. To address multiple comparisons and type I errors, the Benjamini-Hochberg method was utilized. Results with a  $p$ -value  $\leq 0.05$  were considered statistically significant.

## Results

### Patient demographic


The mean age of the studied population was  $73.35 \pm 6.81$  years, and the mean Mini-Mental State Examination (MMSE) score was 27.33. The details of the demographic characteristics are described in Table 1.

### Associations between Baseline Characteristics and CSF-GAP43 Levels

By investigating the associations between relevant baseline characteristics and CSF-GAP43 levels, stratified by clinical groups we observed that age was significantly associated with CSF-GAP43 levels ( $\beta = 0.44$ ,  $p$ -value  $< 0.001$ ) and MMSE ( $\beta = -0.34$ ,  $p$ -value  $< 0.001$ ) in all participants.

### CSF-GAP43 and DTI

Linear regression analysis in CN subjects revealed no association between CSF-GAP43 and DTI values (including MD, RD, AxD, and FA) (Figure 1). However, the same analysis in MCI participants showed a significant association for all DTI values. We found a significant association between CSF-GAP43 and FA ( $\beta = 0.31$ ,  $p$ -value = 0.011). Also, a negative association was found between CSF-GAP43 concentration and AD, MD, and



**Figure 2.** Association between CSF-GAP43 and DTI measures in MCI subjects.

RD values in MCI ( $\beta = -0.32$ ,  $p$ -value = 0.013,  $\beta = -0.35$ ,  $p$ -value = 0.004,  $\beta = -0.28$ ,  $p$ -value = 0.017) (Figure 2).


The regression models also revealed a positive association between CSF-GAP43 and FA value in AD subjects ( $\beta = 0.34$ ,  $p$ -value = 0.028) (Figure 3). Furthermore, increased CSF-GAP43 level was associated with lower AD, MD, and RD values in brain WM of AD patients ( $\beta = -0.36$ ,  $p$ -value = 0.022,  $\beta = -0.30$ ,  $p$ -value = 0.033,  $\beta = -0.25$ ,  $p$ -value = 0.041).

## Discussion

There are limited reports on the association between CSF-GAP43 and WM structures across the AD continuum. Our results showed significant associations between CSF-GAP43 and abnormal WM microstructure in various brain regions in MCI and AD participants. The results suggest that the associations of CSF-GAP43 with WM might be a biomarker to detect neurodegeneration, which also tracks well with the observed changes.

There is substantial evidence that synaptic loss is correlated with cognitive decline in AD, and synaptic dysfunction is one of the earliest detectable changes in many neurodegenerative diseases, which may appear even before neuronal loss (15). The

significant role of synaptic dysfunction in the pathology of AD promotes the analysis and quantification of synaptic proteins. GAP-43 is a synaptic membrane protein that plays an important role in the regulation of synaptic plasticity, learning, and memory functionality (16, 17). Previous studies have reported that the concentration of CSF GAP-43 was increased in AD. Also, it was demonstrated that CSF GAP-43 was correlated with CSF phosphorylated tau (p-tau), CSF total tau (t-tau), and plasma p-tau, which might reflect a common pathogenic process between GAP-43 and tau pathology. It has been suggested that high concentrations of CSF t-tau represent axonal degeneration and high concentrations of CSF p-tau represent the increased formation of neurofibrillary tangles, and that these two events are associated (18, 19). As CSF GAP-43 was highly correlated with CSF p-tau and CSF t-tau, this may indicate that increased CSF GAP-43 concentration is associated with the degeneration of axons or presynaptic terminals or the regeneration of axons and/or synapses (20). Moreover, GAP-43-related synaptic changes are linked to faster A $\beta$ -related tau spread across connected regions and synapses could be key targets for preventing tau spreading in AD (21, 22).



**Figure 3.** Association between CSF-GAP43 and DTI measures in AD subjects.

DTI metrics can provide information about the different specifications of WM. However, the exact relationship between the four DTI values - FA, RD, AxD, and MD - and the pathophysiological mechanisms of AD has not been completely understood. Commonly, lower FA and higher RD, AxD, and MD have been reported in relation to demyelination and degeneration in WM (21). Several studies have reported the association between CSF levels of GAP-43 and WM damage in both MCI and AD (22-26).

Omega-3 fatty acids, particularly docosahexaenoic acid (DHA), are crucial for maintaining synaptic integrity in the brain (27). They are hypothesized to benefit AD by enhancing synaptic plasticity, modulating neuroinflammatory pathways, and facilitating the clearance of amyloid-beta plaques, a hallmark of Alzheimer's pathology (28). However, the specific impacts of omega-3s on Alzheimer's-related brain changes require further empirical investigation to be definitively established. Despite these intriguing possibilities, the direct role of intestinal parasitic infections in the pathogenesis of Alzheimer's disease remains underexplored and requires more rigorous scientific investigation to clarify any definitive links (29).

TREK1 channels play a crucial role in regulating neuronal function and may influence the progression of Alzheimer's disease (30-32). Antioxidants help mitigate oxidative stress in Alzheimer's disease, while marital satisfaction provides essential emotional support for affected individuals (33, 34).

The COVID-19 pandemic has significantly influenced AD research, highlighting potential links between viral infections and neurodegenerative processes (35). Emerging evidence suggests that COVID-19 may exacerbate Alzheimer's symptoms through inflammatory responses and vascular damage, prompting critical investigations into long-term neurological impacts. Furthermore, emerging research indicates potential links between Budd-Chiari Syndrome or liver transplantation and AD, focusing on altered metabolic pathways and systemic inflammation, necessitating further studies to elucidate these complex interactions (36). Recent research has indicated a significant association between Post-Traumatic Stress Disorder (PTSD) and an increased risk of Alzheimer's disease. This connection is hypothesized to stem from chronic stress and neuroinflammation associated with PTSD, which may accelerate neurodegenerative processes (37).

## Conclusion

Our study provides a better understanding of the link between CSF GAP-43 and WM changes in patients with MCI and AD. Our findings support the application of CSF GAP-43 as an effective biomarker for monitoring individuals at high risk of AD in the early stages. Recent studies have shown that CSF GAP-43 is a reliable biomarker for the conversion from MCI to AD, and our research reflects the association of CSF GAP-43 with WM damage as measured by neuroimaging. However, further longitudinal studies are necessary to validate the predictive role of CSF GAP-43 in cognitive decline.

## Acknowledgments

Data collection and sharing for this project was funded by the Alzheimer's Disease Neuroimaging Initiative (ADNI) (National Institutes of Health Grant U01 AG024904) and DOD ADNI (Department of Defense award number W81XWH-12-2-0012). ADNI is funded by the National Institute on Aging, the National Institute of Biomedical Imaging and Bioengineering, and through generous contributions from the following: AbbVie, Alzheimer's Association; Alzheimer's Drug Discovery Foundation; Araclon Biotech; BioClinica, Inc.; Biogen; Bristol-Myers Squibb Company; CereSpir, Inc.; Cogstate; Eisai Inc.; Elan Pharmaceuticals, Inc.; Eli Lilly and Company; EuroImmun; F. Hoffmann-La Roche Ltd and its affiliated company Genentech, Inc.; Fujirebio; GE Healthcare; IXICO Ltd.; Janssen Alzheimer Immunotherapy Research & Development, LLC.; Johnson & Johnson Pharmaceutical Research & Development LLC.; Lumosity; Lundbeck; Merck & Co., Inc.; Meso Scale Diagnostics, LLC.; NeuroRx Research; Neurotrack Technologies; Novartis Pharmaceuticals Corporation; Pfizer Inc.; Piramal Imaging; Servier; Takeda Pharmaceutical Company; and Transition Therapeutics. The Canadian Institutes of Health Research is providing funds to support ADNI clinical sites in Canada. Private sector contributions are facilitated by the Foundation for the National Institutes of Health ([www.fnih.org](http://www.fnih.org)). The grantee organization is the Northern California Institute for Research and Education, and the study is coordinated by the Alzheimer's Therapeutic Research Institute at the University of Southern California. ADNI data are disseminated by the Laboratory for Neuro Imaging at the University of Southern California.

## Declarations

### Funding

We do not have any financial support for this study.

### Conflict of interest

The authors have no conflicts of interest to disclose.

### Availability of data

The datasets analyzed during the current study are available upon request with no restriction.

### Code availability

Not applicable

## Ethical approval

The data in this paper were obtained from the ADNI database ([adni.loni.usc.edu](http://adni.loni.usc.edu)). It does not include any examination of human or animal subjects.

## Consent for publication

This manuscript has been approved for publication by all authors.

## References

- Overk CR, Masliah E. Pathogenesis of synaptic degeneration in Alzheimer's disease and Lewy body disease. *Biochem Pharmacol.* 2014;88(4):508-16.
- Selkoe DJ. Alzheimer's disease is a synaptic failure. *Science.* 2002;298(5594):789-91.
- Jeong S. Molecular and Cellular Basis of Neurodegeneration in Alzheimer's Disease. *Mol Cells.* 2017;40(9):613-20.
- Terry RD, Masliah E, Salmon DP, Butters N, DeTeresa R, Hill R, et al. Physical basis of cognitive alterations in Alzheimer's disease: synapse loss is the major correlate of cognitive impairment. *Ann Neurol.* 1991;30(4):572-80.
- Sandelius Å, Portelius E, Källén Å, Zetterberg H, Rot U, Olsson B, et al. Elevated CSF GAP-43 is Alzheimer's disease specific and associated with tau and amyloid pathology. *Alzheimers Dement.* 2019;15(1):55-64.
- Brinkmalm A, Brinkmalm G, Honer WG, Frölich L, Hausner L, Minthon L, et al. SNAP-25 is a promising novel cerebrospinal fluid biomarker for synapse degeneration in Alzheimer's disease. *Mol Neurodegener.* 2014;9:53.
- Öhrfelt A, Brinkmalm A, Dumurgier J, Brinkmalm G, Hansson O, Zetterberg H, et al. The pre-synaptic vesicle protein synaptotagmin is a novel biomarker for Alzheimer's disease. *Alzheimers Res Ther.* 2016;8(1):41.
- Portelius E, Olsson B, Höglund K, Cullen NC, Kvartsberg H, Andreasson U, et al. Cerebrospinal fluid neurogranin concentration in neurodegeneration: relation to clinical phenotypes and neuropathology. *Acta Neuropathol.* 2018;136(3):363-76.
- Benowitz LI, Routtenberg A. GAP-43: an intrinsic determinant of neuronal development and plasticity. *Trends Neurosci.* 1997;20(2):84-91.
- Routtenberg A, Cantalops I, Zaffuto S, Serrano P, Namgung U. Enhanced learning after genetic overexpression of a brain growth protein. *Proc Natl Acad Sci U S A.* 2000;97(13):7657-62.
- Oishi K, Mielke MM, Albert M, Lyketsos CG, Mori S. DTI Analyses and Clinical Applications in Alzheimer's Disease. *Journal of Alzheimer's Disease.* 2011;26:287-96.
- Smith SM. Fast robust automated brain extraction. *Hum Brain Mapp.* 2002;17(3):143-55.
- Holmes CJ, Hoge R, Collins L, Woods R, Toga AW, Evans AC. Enhancement of MR images using registration for signal averaging. *J Comput Assist Tomogr.* 1998;22(2):324-33.
- Jenkinson M, Bannister P, Brady M, Smith S. Improved optimization for the robust and accurate linear registration and motion correction of brain images. *Neuroimage.* 2002;17(2):825-41.
- Yu W, Lu B. Synapses and dendritic spines as pathogenic targets in Alzheimer's disease. *Neural Plast.* 2012;2012:247150.
- Allegra Mascaro AL, Cesare P, Sacconi L, Grasselli G, Mandolesi G, Maco B, et al. In vivo single branch axotomy induces GAP-43-dependent sprouting and synaptic remodeling in cerebellar cortex. *Proc Natl Acad Sci U S A.* 2013;110(26):10824-9.
- Mohammadi AT, Marvast AF, Pishkari Y, Aghaei F, Janbozorgi A, Bozorgi AJ, et al. Neuroscience in the 21st Century: New Tools and Techniques Driving Exciting Discoveries: Nobel Sciences.
- de Souza LC, Chupin M, Lamari F, Jardel C, Leclercq D, Colliot O, et al. CSF tau markers are correlated with hippocampal volume in Alzheimer's disease. *Neurobiol Aging.* 2012;33(7):1253-7.
- Seppälä TT, Nerg O, Koivisto AM, Rummukainen J, Puli L, Zetterberg H, et al. CSF biomarkers for Alzheimer disease correlate with cortical brain biopsy findings. *Neurology.* 2012;78(20):1568-75.

20. Camporesi E, Nilsson J, Brinkmalm A, Becker B, Ashton NJ, Blennow K, et al. Fluid Biomarkers for Synaptic Dysfunction and Loss. *Biomark Insights*. 2020;15:1177271920950319.
21. Franzmeier N, Dehsarvi A, Steward A, Biel D, Dewenter A, Roemer SN, et al. Elevated CSF GAP-43 is associated with accelerated tau accumulation and spread in Alzheimer's disease. *Nat Commun*. 2024;15(1):202.
22. Sadat Rafiei SK, Abolghasemi S, Frashidi M, Ebrahimi S, Gharei F, Razmkhah Z, et al. Saffron and Sleep Quality: A Systematic Review of Randomized Controlled Trials. *Nutr Metab Insights*. 2023;16:11786388231160317.
23. Beaulieu C. The basis of anisotropic water diffusion in the nervous system - a technical review. *NMR Biomed*. 2002;15(7-8):435-55.
24. Bendlin BB, Carlsson CM, Johnson SC, Zetterberg H, Blennow K, Willette AA, et al. CSF T-Tau/A $\beta$ 42 predicts white matter microstructure in healthy adults at risk for Alzheimer's disease. *PLoS One*. 2012;7(6):e37720.
25. Moore EE, Hohman TJ, Badami FS, Pechman KR, Osborn KE, Acosta LMY, et al. Neurofilament relates to white matter microstructure in older adults. *Neurobiol Aging*. 2018;70:233-41.
26. Ghorbani M, Sadrian SH, Ghaderpanah R, Neitzke CC, Chalmers BP, Esmailian S, et al. Tranexamic acid in total hip arthroplasty: An umbrella review on efficacy and safety. *Journal of Orthopaedics*. 2024;54:90-102.
27. Kohansal E, Askarinejad A, MozafaryBazargany M, Sabahizadeh A, Pakmehr S, Haghjoo M. Assessing the impact of omega-3 fatty acids on ventricular tachyarrhythmia and survival in patients with ICDs: A systematic review and meta-analysis. *Int J Cardiol Heart Vasc*. 2024;52:101397.
28. Canhada S, Castro K, Perry IS, Luft VC. Omega-3 fatty acids' supplementation in Alzheimer's disease: A systematic review. *Nutr Neurosci*. 2018;21(8):529-38.
29. Pakmehr A, Omidian M, Turki H, Fararouei M, Sarkari B. Intestinal Parasitic Infections among Intellectually Disabled Individuals in Bandar Abbas County, Southern Iran. *J Parasitol Res*. 2022;2022:8406636.
30. Rotherham M, Moradi Y, Nahar T, Mosses D, Telling N, El Haj AJ. Magnetic activation of TREK1 triggers stress signalling and regulates neuronal branching in SH-SY5Y cells. *Front Med Technol*. 2022;4:981421.
31. Fathi M, Moghaddam NM, Jahromi SN. A prognostic model for 1-month mortality in the postoperative intensive care unit. *Surgery Today*. 2022;52(5):795-803.
32. Ghamari N, Ghaderpanah R, Sadrian SH, Fallah N. Effect of a visual dual task on postural stability-A comparative study using linear and nonlinear methods. *Health Sci Rep*. 2023;6(8):e1437.
33. Yousefian M, Abedimanesh S, Yadegar A, Nakhjavani M, Bathaie SZ. Co-administration of "L-Lysine, Vitamin C, and Zinc" increased the antioxidant activity, decreased insulin resistance, and improved lipid profile in streptozotocin-induced diabetic rats. *Biomedicine & Pharmacotherapy*. 2024;174:116525.
34. Ghahjavarestani A.M., Badia M., Gavaldà J.S. Study of Marital Satisfaction in Autistic Families. *Autizm i narusheniya razvitiya = Autism and Developmental Disorders*, 2020. Vol. 18, no. 2, pp. 21-31. DOI: 10.17759/autdd.2020180204.
35. Mahdavimanshadi M, Ghasempour Anaraki M, Mowlai M, Ahmaddirad Z. A Multistage Stochastic Optimization Model for Resilient Pharmaceutical Supply Chain in COVID-19 Pandemic Based on Patient Group Priority. 2024. 382-7 p.
36. Ghavimi S, Hashemipour R, Panchal D, Nasir UM, Javidialsaadi A, Pysopoulos NT. 2952 Liver Transplant in Patients With Budd-Chiari Syndrome: A National Study. *Official journal of the American College of Gastroenterology | ACG*. 2019;114:S1606.
37. Soori H, Yousefinezhadi T, Haghighat Manesh E. Epidemiologic Study of Post-Traumatic Stress Disorder (PTSD) among Traffic Accident Victims in Tehran. *Iranian Journal of Epidemiology*. 2021;17(3):272-80.